

8/148/60/000/002/004/008

Corrosion Resistance of Austenite Steels After Pressure Working at Low Temperatures

the steel was less impaired and a maximum on the curve ("weight loss versus degree of compression") was not observed. After rolling at low temperature X-ray examinations of the phase composition in austenite steels were carried out to determine, to which amount of martensite corresponded the maximum decrease of corrosion resistance. Comparison of curves (Figures 2, 3) show a maximum weight loss in 1Kh18N9 steel after compression by 15 - 20%, which corresponded to about 50% martensite. If compression was increased up to 40%, corrosion resistance improved and then became almost equal to that of steel rolled at room temperature. Thus, after rolling at -183°C and 40% compression, the weight loss was 14 g/m^2 and after rolling at room temperature it was 11 g/m^2 . Improved corrosion resistance in the presence of a martensite content of over 50% in steel rolled at low temperatures, proved that high-alloy martensite ensured sufficiently high corrosion resistance. The conclusion is drawn that in austenite steels with non-stable austenite,

Card 2/3

S/148/60/000/002/004/004

Corrosion Resistance of Austenitic Steels After Pressure Working at Low
Temperatures

the pressure working at low temperatures, carried out to raise the strength,
dit not considerably reduce their corrosion resistance.
There are: 3 graphs and 4 references, 3 of which are Soviet and 1 English

ASSOCIATION Khar'kovskiy aviatsionnyy institut (Khar'kov Aviation Institute)

SUBMITTED: November 21, 1958

Card 3/3

PISARENKOVICH, I.N., inzhener.

Over-all mechanisation in building protective hydrotechnical
structures. Stroi. prom. 35 no.3:18-24 Mr '57. (MLRA 10:4)
(Dnieper River--Flood dams and reservoirs)
(Building machinery)

PISAREVA, M. G.

PISAREVA, M. G. -- "Methods of Effective Fertilization of Annual Lupine in Combination with Peat and Straw." Moscow, 1956. (Dissertation for the Degree of Candidate in Agriculture Sciences).

So.: Knizhnaya Litopis', No. 7, 1956.

PISAREVA, N.N.

Fraction-quadratic integral of geodesic lines in m -dimensional
spaces of affine connectivity. Dokl.AN SSSR 108 no.2:198-200
My '56.
(MIRA 9:9)

1.Gor'kovskiy gosudarstvennyy universitet imeni N.I.Lobachevskogo.
Predstavleno akademikom I.G.Petrovskim.
(Spaces, Generalized)

SUBJECT USSR/MATHEMATICS/Geometry CARD 1/3 PG - 104
 AUTHOR PISAREVA N.M.
 TITLE On the rational-quadratic integral of the geodesic lines in spaces of affine connection.
 PERIODICAL Mat. Sbornik, n. Ser. 36, 169-200 (1955)
 reviewed 6/1956

In the first part of the paper the two-dimensional spaces A_2 with affine connection are considered. The investigation bases essentially on the theory of the nets of non-metric spaces of affine connection given by Norden. Some theorems on the relations between rational quadratic integrals and different nets are given; e.g.: In order that

$$(1) \quad \frac{a_{ij} du^i du^j}{b_{ij} du^i du^j} = C$$

is an integral of the geodesic lines of the A_2 , it is necessary and sufficient that the tensors a_{ij} , b_{ij} and any linear combination of both represent geodesic nets in the A_2 . The geodesic nets $a_{ij} du^i du^j = 0$ and $b_{ij} du^i du^j = 0$ are denoted as basis nets of the integral (1). In order that the geodesic lines of the A_2 possess a first rational linear integral it is necessary and sufficient that in A_2 there exist two geodesic nets the Chebyev vectors of which differ by the gradient of a function. These nets then are basis nets of the integral.

Mat. Sbornik, n. Ser. 36, 169-200 (1955)

CARD 2/3

PG - 104

Nets with the same χ Coby vectors are called special nets. In order that (1) is an integral of the geodesic lines, where the ratio of the discriminants of a_{ij} and b_{ij} is constant, a_{ij} and b_{ij} must represent special nets which then are basis nets too. Then spaces of affine connection are sought the geodesic lines of which possess first integrals of the form (1). A generalization of the Darboux theorem on the geodesic mapping is proved: In order that the Weyl's space W_2 admits a geodesic gradient mapping onto another W_2 , it is necessary and sufficient that the geodesic lines of the W_2 possess an integral (1), one basis net of which is an isotropic net of the given space W_2 . Three special cases of the A_2 which admit (1) are considered.

The second part of the paper relates to more-dimensional spaces of affine connection. As more-dimensional analogue of the quasi-euclidean spaces, reducible Weyl spaces are introduced, i.e. such Weyl spaces in which there exists a tensor b_{ij} being different from the measure tensor, which for a certain vector N_k satisfies the condition: $b_{ij,k} = b_{ij}N_k$. It is remarked that the reducible Weyl spaces with two one to another orthogonal systems of completely geodesic surfaces are no Riemannian spaces. For more than two systems of the above kind they are Riemannian spaces. After these preliminary remarks the results of the first part are generalized to the more-dimensional case. As an analogue of the geodesic net in A_2 , in the A_n the notion of the geodesic conic field due to Šapiro (Doklady Akad. Nauk 39, 6-10 (1943)) is used. Some

Mat. Sbornik, n. Ser. 36, 169-200 (1955)

CARD 3/3

PG - 104

results of the first part then can be transferred completely: In order that (1) is the integral of the geodesic lines of the Δ_n , it is necessary and sufficient that the tensors a_{ij} and b_{ij} and any linear combination of it represent geodesic conic fields of the Δ_n . For the generalization of the results on special geodesic nets the notion of a projective-metric conic field is introduced. The tensor of this field satisfies the condition

$$na_{ij,k} = \frac{n}{2} (t_1 a_{kj} + t_j a_{ki}) + (a_{\alpha\beta k} \tilde{a}^{\alpha\beta} - t_k) a_{ij},$$

where t_1 is the Cebyshev vector of the conic field. Two conic fields with the same t_1 are called special conic fields. By introduction of these new notions the remaining results of the first part can be transferred to the more-dimensional case.

INSTITUTION: Gorki.

1940-1941.

Disanova, I. "The role of the Soviet Union in the development of the world economy," in: "The role of the Soviet Union in the development of the world economy," (Moscow, 1941), p. 100.

3. The role of the Soviet Union in the development of the world economy, (Moscow, 1941), p. 100.

ANOSOV, N.N; PISAREVA, N.A. (Leningrad)

Therapy of neuralgic syndrome of various origins with phenadon.
6-dimethylamino-4,4-diphenyl -3-heptanone HCl. Sov. med. 18 no.11:
11-12 N '54. (MLRA 7:12)

(ANALGESICS, ther. use

6-dimethylamino-4,4-diphenyl -3-heptanone HCl in
neuralgia)

(NEURALGIA, therapy

6-dimethylamino-4,4-diphenyl -3-heptanone HCl)

AUTHOR: Pisareva, N. M. (Gor'kiy). 190

TITLE: Weyl spaces containing a projective system of paths.
(Prostranstva Veylya, vklyuchayushchie proyektivnuyu
sistemu putey).

PERIODICAL: "Matematicheskiiy Sbornik" (Mathematical Symposium),
1957, Vol.41(83), No.2, pp.231-238 (U.S.S.R.)

ABSTRACT: The object of this paper is to review Weyl spaces
which "include" a projective system of paths and to
establish a relation between these spaces and reducible
Weyl spaces. It appears that both these classes of
spaces have a general subclass of spaces characterised
in the following manner:-
A space of the subclass contains two mutually
orthogonal systems of fully geodesic surfaces and one
of these systems defines a geodesic field of m-dimensional
directions in the space under investigation.
Consider a space of affine connectivity of $n + m$
dimensions A_{n+m} . m linearly independent vector fields
 V^a define in this space a certain field of m-dimensional
directions. The m-dimensional field is called holonomic
if the following conditions are satisfied:-
(1) Through each point A_{n+m} passes one of the
spaces of the system, a
(2) The vectors V^a defining the direction of the
field at a given point are tangential to the corresponding

Weyl spaces containing a projective system of paths. 199
(Cont.)

subspace of the system. A holonomic field of m -dimensional directions is called geodesic if the manifolds of $m + 1$ dimensions formed the above subspaces passing through a point of any geodesic line of A_{n+m} are fully geodesic.

The conditions (14), (15) (see p.235) and (18) (p.236), are sufficient for the spaces W_{n+m} to include a projective system of paths. The geodesic projection of the space W_{n+m} by means of the coordinates of the manifolds $v^i = \text{const.}$ ($i = 1, 2, \dots, n$) on the manifolds $v^a = \text{const.}$ is a system of paths, ($a = n + 1, \dots, n + m$).

If it is supposed that the n -dimensional manifolds (8) (p.234) are fully geodesic, then the space W_{n+m} would be a reducible Weyl space, but not a reducible space of the same general type because the $(m + 1)$ -dimensional manifolds of subspaces mentioned above, passing through points of an arbitrary geodesic line of the space W_{n+m} are fully geodesic. There are three references, all Russian.

Submitted 19/1/56.

PISAREVA, N.M.

PISAREVA, N.M. (Gor'kiy)

Fractional-quadratic integral of geodesic lines of affinely connected spaces. Mat.sbor. 36 no.1:169-200 Ja-P '55. (MLRA 8:2)
(Integrals) (Spaces, Generalized)(Line geometry)

LYULICHEVA, N.N.; PISAREVA, N.V.

Heat treatment of cold-worked Kh18N9T steel. Metalloved. i term.
obr. met. no.10:41-42 0 '63. (MIRA 16:10)

1. Khar'kovskiy aviatsionnyy institut.

8/187/39/000/00/030/030
8031/8013

AUTHOR: Galstabin, V.I.

TITLE: The Scientific-Technical Conference at Kharkov Aviation Institute

PERIODICAL: Izvestiya Vysheishykh uchebnykh zavedeniy, Aviatsionnaya tekhnika, 1959, No. 8, pp 161-165 (USSR)

ABSTRACT: In May 1959, the 16th Conference of Professors and Teaching Staff took place.

Card 8/11 The Technology of Aircraft Construction and Metal Working Institute. "The Model of the Plasticity of Metals" by Galstabin, V.I.

Card 9/11 "On the Problem of Constructing Second Order Curves in Components from Sheet Metals" by Galstabin, V.I.

Card 10/11 "On the Problem of Constructing Second Order Curves in Components from Sheet Metals" by Galstabin, V.I.

Card 11/11 "On the Problem of Constructing Second Order Curves in Components from Sheet Metals" by Galstabin, V.I.

Card 12/11 "On the Problem of Constructing Second Order Curves in Components from Sheet Metals" by Galstabin, V.I.

Card 13/11 "On the Problem of Constructing Second Order Curves in Components from Sheet Metals" by Galstabin, V.I.

Card 14/11 "On the Problem of Constructing Second Order Curves in Components from Sheet Metals" by Galstabin, V.I.

Card 15/11 "On the Problem of Constructing Second Order Curves in Components from Sheet Metals" by Galstabin, V.I.

Card 16/11 "On the Problem of Constructing Second Order Curves in Components from Sheet Metals" by Galstabin, V.I.

Card 17/11 "On the Problem of Constructing Second Order Curves in Components from Sheet Metals" by Galstabin, V.I.

Card 18/11 "On the Problem of Constructing Second Order Curves in Components from Sheet Metals" by Galstabin, V.I.

Card 19/11 "On the Problem of Constructing Second Order Curves in Components from Sheet Metals" by Galstabin, V.I.

Card 20/11 "On the Problem of Constructing Second Order Curves in Components from Sheet Metals" by Galstabin, V.I.

Card 21/11 "On the Problem of Constructing Second Order Curves in Components from Sheet Metals" by Galstabin, V.I.

Card 22/11 "On the Problem of Constructing Second Order Curves in Components from Sheet Metals" by Galstabin, V.I.

Card 23/11 "On the Problem of Constructing Second Order Curves in Components from Sheet Metals" by Galstabin, V.I.

Card 24/11 "On the Problem of Constructing Second Order Curves in Components from Sheet Metals" by Galstabin, V.I.

Card 25/11 "On the Problem of Constructing Second Order Curves in Components from Sheet Metals" by Galstabin, V.I.

Card 26/11 "On the Problem of Constructing Second Order Curves in Components from Sheet Metals" by Galstabin, V.I.

Card 27/11 "On the Problem of Constructing Second Order Curves in Components from Sheet Metals" by Galstabin, V.I.

Card 28/11 "On the Problem of Constructing Second Order Curves in Components from Sheet Metals" by Galstabin, V.I.

Card 29/11 "On the Problem of Constructing Second Order Curves in Components from Sheet Metals" by Galstabin, V.I.

Card 30/11 "On the Problem of Constructing Second Order Curves in Components from Sheet Metals" by Galstabin, V.I.

Card 31/11 "On the Problem of Constructing Second Order Curves in Components from Sheet Metals" by Galstabin, V.I.

Card 32/11 "On the Problem of Constructing Second Order Curves in Components from Sheet Metals" by Galstabin, V.I.

Card 33/11 "On the Problem of Constructing Second Order Curves in Components from Sheet Metals" by Galstabin, V.I.

Card 34/11 "On the Problem of Constructing Second Order Curves in Components from Sheet Metals" by Galstabin, V.I.

Card 35/11 "On the Problem of Constructing Second Order Curves in Components from Sheet Metals" by Galstabin, V.I.

Card 36/11 "On the Problem of Constructing Second Order Curves in Components from Sheet Metals" by Galstabin, V.I.

Card 37/11 "On the Problem of Constructing Second Order Curves in Components from Sheet Metals" by Galstabin, V.I.

Card 38/11 "On the Problem of Constructing Second Order Curves in Components from Sheet Metals" by Galstabin, V.I.

Card 39/11 "On the Problem of Constructing Second Order Curves in Components from Sheet Metals" by Galstabin, V.I.

Card 40/11 "On the Problem of Constructing Second Order Curves in Components from Sheet Metals" by Galstabin, V.I.

Card 41/11 "On the Problem of Constructing Second Order Curves in Components from Sheet Metals" by Galstabin, V.I.

LYULICHEVA, N.N.; PISAREVA, N.V.

Corrosion resistance of austenitic steels following press working
at low temperatures. Izv. vys. ucheb. zav.; Chern. met. no. 1:
78-80 '60. (MIRA 15:5)

1. Khar'kovskiy aviatsionnyy institut.
(Steel, Stainless--Corrosion)
(Metals at low temperatures)

18(2)

AUTHORS: Lyulicheva, N. N., Pisareva, N. V.

SOV, 163-51-2-1 41

TITLE: The Mechanical Properties of Cold Hardened Chromium-nickel-austenite Steel of the Type 18-8 at Low Temperatures
(Mekhanicheskiye svoystva nagartovannykh khromonikelevykh austenitnykh staley tipa 18-8 pri nizkikh temperaturakh)

PERIODICAL: Nauchnyye doklady vysshey shkoly. Metallurgiya, 1961, Nr 2, pp 217-220 (USSR)

ABSTRACT: The mechanical properties of stainless austenite steels of the type 18Cr13Ni and 18Cr13Ni were investigated at temperatures of +20 and -195° after rolling at room temperature (Fig 1). The dependence of the relative extension of the austenite steel of the type 18-8 on different temperatures is given in figure 2. The higher plasticity of the austenite steels determined at 183° after rolling at +20° is caused by the occurrence of cubic face-centered lattices. The mechanical properties of the metals and alloys after the treatment under pressure at room temperature are summarized in the table. The transformation of martensite into austenite steel during the deformation process at low temperature increases the strength of the alloys. A previous cold hardening

Card 1/2

SOV, 1958, 1-1-1, 4
The Mechanical Properties of Cold Hardened Chromium-nickel-austenitic
Steel of the Type 18-8 at Low Temperatures

is excellent for the use of austenitic steel for low temperatures since the flow limit is thus increased without reduction of the plasticity. There are 2 figures, 1 table, and 4 references, 4 of which are Soviet.

ASSOCIATION: Khar'kovskiy aviatsionnyy institut
(Khar'kov Institute of Aviation)

SUBMITTED: May 28, 1958

Card 2/2

SOKOLOVA, Ye.B.; SHUBANOVA, M.I.; CHZHOU KHEH-TSZIN' [Chou Heng-chin];
PISARENKO, S.A.

Use of fulvenes for the synthesis of homologs and analogs of
ferrocene. Zhur. ob. khim. 34 no.8:2693-2696 Apr 1964.
(MIRA 17:9)

KOLOSOVA, N.N.; PISAREVA, T.N. (Leningrad)

Multiple lipomas. Vop. neirokhir. 27 no.2:53-54 Mr-Apr '63.
(MIRA 17:2)

1. Kafedra nervnykh bolezney (zav. - prof. Ye.F. Davidenkova)
i kafedra patologicheskoy anatomii (zav. - prof. V.G.
Chudakov) Leningradskogo pediatricheskogo meditsinskogo
instituta.

LANDA, Ya.Kh., dotsent; PISAREVA, T.N., kand.med.nauk

Periarteritis nodosa; according to data from the V.V. Knibyshev
Hospital. Vop.pat.krovi i krovoobr. no.6:229-235 '61.

(PERIARTERITIS NODOSA)

(MIRA 16:3)

PISAREVA, T. N., Cand of Med Sci -- (diss) "Circulatory Disturbances and Brain Hemorrhages in New-born Children," Leningrad, 1959, 13 pp (Leningrad Pediatrics Institute) (KL, 7-60, 110)

PISARENVA, V.V.

Diffraction of radiowaves on random nonuniformities and intensity variations of solar and cosmic radiowave emission [with summary in English]. Astron. zhur. 35 no.1:112-128 Ja-F '58. (MIRA 11:3)

1. Gor'kovskiy gosudarstvennyy universitet.
(Solar radiation) (Diffraction) (Radio astronomy)

1(1)
 AUTHOR. Pisareva, V.V. SOV. ASTRON. 1959, 16, 3, 427-433
 TITLE On the Variations of the Intensity of Radio Emission Scattered on Corona Nonuniformities
 PERIODICAL Astronomi beskiy zhurnal, 1959, Vol. 16, Nr 3, pp 427-433 (USSR)
 ABSTRACT For the investigation of the intensity fluctuations of the solar radio emission from September 15 to November 1, 1957 in Leningrad and Irkutsk simultaneous observations were carried out (the towns have nearly the same latitude and are at a distance of ~5000 km from each other). It was conjectured that in this manner a timely shift of the diffraction image of ~10 sec can be asserted; that would confirm the assumption that a part of the fluctuations of solar radio emission can be attributed to the diffraction of the radiation at the coronal nonuniformities. But the conjecture was not confirmed. It is shown that for the confirmation of the mentioned shift a 10 times greater antenna surface (100 m²) and a more sensible apparatus would be necessary. As another possibility for the investigation of the nonuniformities of the corona the author recommends the measurement of the intensity of the radio emission of the Crab nebula, when this nebula is covered by the solar corona. It is shown that in

Card 1/2

5

On the Variations of the Intensity of Radio Emission, SOV/33-36-1 5/2.
Scattered on Corona. Nonuniformities

a large distance from the solar center the apparent increase in the size of the source by scattering at the coronal rays cannot be explained. It is assumed that the corona has also smaller nonuniformities. The author thanks Professor V.L.Ginzburg and G.G. Getmantsev for advice. There is 1 figure, and 12 references, 7 of which are Soviet, 3 English, 1 Dutch, and 1 American.

ASSOCIATION: Radiofizicheskiy institut pri Gorkovskoye gosudarstvennoy universite-
imeni V.I.Lobachevskogo (Radiophysical Institute at the Gorky State
University imeni V.I.Lobachevskogo)

SUBMITTED: July 15, 1958

Card 2/2

PISAREVA, V.V.

Applicability range of the "smooth" perturbation method in the
problem of wave propagation in an irregular medium. Akust.zhur.
87-91 '60.

(MIRA 14:5)

1. Gor'kovskiy gosudarstvennyy universitet.
(Sound—Transmission)

PISAREVA, V.V.

Polarization of nonthermal Galactic radio emission of radio emission
from the Crab nebula. Izv.vys.ucheb.zav.; radiofiz. 3 no.2:
165-179 '60. (MIRA 13:7)

1. Nauchno-issledovatel'skiy radiofizicheskiy institut pri
Gor'kovskom universitete.
(Cosmic radiations, Radio-frequency)

PISAREVA, V. V.

Cand Phys-Math Sci - (diss) "Statistical theory of the propagation of radio waves in the ionosphere and under conditions of space." Gor'kiy, 1961. 8 pp including cover; (Ministry of Higher and Secondary Specialist Education RSFSR, Gor'kiy State Univ imeni N. I. Lobachevskiy, Scientific Research Radiophysics Inst "NIRFI"); 200 copies; price not given; bibliography at end of text (17 entries); (KL, 6-61 sup, 195)

9.9110

S/141/61/004/002/016/017
E032/E114

AUTHOR: Pisareva, V V

TITLE: On the limits of applicability of the method of continuous perturbations as applied to the propagation of radiation through a medium with irregularities

PERIODICAL: Izvestiya vysshikh uchebnykh zavedeniy, Radiofizika 1961 Vol.4. No.2. pp. 376-377

TEXT: The limits of applicability of the method of continuous perturbations were discussed in a previous paper by the present author (Ref.1: Akust. zh. Vol.6, 87 (1960)). It is stated that the method used in that paper is not sufficiently correct. However more rigorous analysis of the problem, briefly described in the present note, leads to exactly the same results as those given in Ref.1. In particular it is shown that, when $D \ll 1$.

$$\frac{\alpha^2 |\psi_2|}{\alpha \sqrt{\psi_1^2}} = \sqrt{\frac{\sqrt{\pi} L^3 \mu^2}{i^3}} \approx 0.6 \sqrt{[\ln(A/A_0)]^2} \ll 1$$

Card 1/3

On the limits of applicability of S/141/61/004/002/016/017
and EO32/E114

$$\frac{a^2 |\psi_2|}{a \sqrt{\psi_1^2}} = \sqrt{\frac{\sqrt{\pi}}{2} k_0^2 \ell L \mu^2} \approx 0.7 \sqrt{(a \psi_1)^2} \ll 1$$

when $D \gg 1$. In these expressions $D = 4L/k_0 \ell^2$, L is the range, k_0 is the wave number in vacuum, ℓ is the average dimension of the irregularities in the medium, $a = \sqrt{\mu^2}$, μ is the average variation in the refractive index (measured from the mean) and $a\psi_1$ and $a^2\psi_2$ are the first terms in the expansion for ψ in powers of the small parameter μ . Furthermore A and A_0 are the amplitudes of the perturbed and unperturbed waves (cf. Ref. 1). It is argued that the condition for the applicability of the method of continuous perturbations, namely

$$\sqrt{[\ln(A_1/A_0)]^2} \ll 1, \quad D \ll 1,$$

is necessary and sufficient:

Card 2/3

On the limits of applicability of ... S/141/61/004/002/016/017
E032/E114

There are 4 references: 3 Soviet and 1 English. The English
language reference reads as follows:
Ref 4: I.J. Howells Phil. Trans of the Royal Soc. of London.
Vol.252, 431 (1960)

ASSOCIATION: Nauchno-issledovatel'skiy radiofizicheskiy institut
pri Gor'kovskom universitete
(Scientific Research Institute of Radiophysics at
Gor'kiy University)

SUBMITTED: December 17 1960

Card 3/3

PISAREVA, V.V.

Limits of applicability of the method of smooth perturbations
in the problem of the propagation of radiation through a medium
containing inhomogeneities. Akust. zhur. 7 no.3:387 '61. (MIRA 14:9)

1. Nauchno-issledovatel'skiy Radiofizicheskiy institut ;ri
Gor'kovskom gosudarstvennom universitete.

(Boundary value problems) (Ratiation)

GINZBURG, V.L.; PISAREVA, V.V.

Polarization of the radio emission from discrete sources, and the
exploration of the metagalactic and galactic spaces and spaces
near the sun. Izv. vys. ucheb. zav.; radiofiz. 6 no.5:877-888
'63. (MIRA 16:12)

1. Nauchno-issledovatel'skiy radiofizicheskiy institut pri
Gor'kovskom universitete.

KOROBKOV, Yu.S.; PISAREVA, V.V.

Latitudinal distribution of the ionospheric absorption of a short
radio-frequency radiation. Geomag. i aer. 5 no.1:173-177 Ja-F '65.
(MIRA 38:4)

1. Nauchno-issledovatel'skiy radiofizicheskiy institut pri
Gor'kovskom universitete.

L 52187-65 EMT(a)/FSS-2/EMT(1)/EEC(a)/EEC(j)/EEC(k)-2/EEC(f)/EEC(r)/EWG(v)/
 FCC/EEC-L/EEC(t)/EEC(c)-2/EWA(h) Fm-L/Po-L/Pp-L/Pe-5/Pq-L/Pac-L/Pae-2/PeB/Pi-L
 ACCESSION NR: AP5014102 AST/GW UR/0203/65/005/003/0423/0428
 550.385

AUTHORS: Korobkov, Yu. S.; Pisareva, V. V.

TITLE: Study of the inhomogeneities of electron concentration in the ionosphere in the region of the Tycho ocean with the help of artificial earth satellite signals

SOURCE: Geomagnetizm i aeronomiya, v. 5, no. 3, 1965, 423-428

TOPIC TAGS: electron, ionosphere, artificial satellite, antenna/ Explorer VII satellite, N 10 oscillograph

ABSTRACT: The longitudinal dependence of the index of fluctuation of the signal from the artificial satellite Explorer-VII in the range of latitudes from 41°N to 18°S was studied. The correlation between the index of fluctuation, the diffuseness of the F-2 layer, and the frequencies of occurrence of E_s has been established. The diurnal dependence of the index of fluctuation and its relation to the magnetic activity are briefly considered and the magnitudes of the inhomogeneities are estimated. The frequency of the signal from the satellite was 19 992 megacycles. The authors were interested only in observing signals of relatively large wavelengths, $\lambda \sim 15$ m. The signal was received by half-wave dipole antennae at an altitude of 0.5 m above the deck of the vessel. After detection, the signal was sent through an

L 52187-65

ACCESSION NR: AP5014102

integrating network having a time constant of 0.02 sec, and was registered on an N-10 oscillograph. The data about the ionosphere were obtained in an AIS station. The diffraction pattern on the earth's surface was obtained, and the following formula was used for τ , the time for irregular changes in the signal in the Fraunhofer region

$$\tau = \frac{\Delta \xi}{v \sqrt{(\Delta \varphi)^2}} \frac{h_1}{h_2} \cdot (h = h_1 h_2 / (h_1 + h_2)).$$

Here $\Delta \xi$ is the size of the inhomogeneity, v the velocity of the satellite, $\Delta \varphi$ the phase shift due to the inhomogeneity, h_1 the altitude of the inhomogeneity, and h_2 the altitude of the satellite. It was found from these observations that the index of fluctuation at night did not have a maximum in time. In most cases it was found to be well correlated to the diffuseness of the F-2 layer. The coefficient of correlation was found to be 0.16. Orig. art. has: 3 figures and 2 formulas.

ASSOCIATION: Nauchno-issledovatel'skiy radiofizicheskiy institut pri Gorkovskom universitete (Radiophysical Scientific Research Institute, Gorkiy University)

SUBMITTED: 15Jul64

ENCL: 00

SUB CODE: AA, BV

NO REF SOV: 002

OTHER: 008

c.28/2/2

L 11369-65

GN/WS

ENT(1)/ENG(v)/FOC/EEC-4/EEC(t)/EWA(h)

Po-4/Pe-5/Pq-4/Pae-2/Peb/P1-4

ACCESSION NR: AP4046283

S/0203/64/004/005/0066/0872

AUTHOR: Korobkov, Yu. S.; Pisareva, V. V.

TITLE: Several results of investigations of E_s layers in the region of the Pacific B

SOURCE: Geomagnetizm i aeronomiya, v. 4, no. 5, 1964, 866-872

TOPIC TAGS: ionospheric sporadic layer, appearance frequency, local time, E_s layer type, semitransparency

ABSTRACT: Investigations of ionospheric sporadic E_s layers were carried out over the Pacific between the latitude parallels 42°N and 18°S in the summer of 1962. The whole region was divided into three zones — zone I, equatorial; zone II, northern tropical; and zone III, northern subtropical. E_s types c, i, f, s, and q were analyzed. The maximum appearance frequency of the f-type took place between 2 and 3 p.m. local time. The c-type had a minimum appearance at noon. The i-type had a minimum at 2 p.m. These charac-

Card 1/2

L 11369-65

ACCESSION NR: AP4046283

Characteristics relate to zone I. With increased latitude, the maximum of the appearance frequency is transferred to the night hours. The maximum appearance frequency of the I-type takes place in zone II. A maximum reflection from the I-type was observed near the equator. Some of the E_s types totally screen the reflection of upper ionospheric layers, and others are semitransparent, depending upon their structure. The f-type is opaque at night. Orig. art. has: 5 figures.

ASSOCIATION: Radiofizicheskiy institut pri Gor'kovskom gosudarstvennom universitete (Radiophysical Institute at Gorkiy State University)

SUBMITTED: 08Jan64

ATD PRESS: 3114

ENCL: 000

SUB CODE: AA.01

NO. REF SOVI: 003

OTHER: 003

Cord 2/2

KARLIN, M.I., kand.med.nauk; LIPSKAYA, M.I., kand.med.nauk; PISAREVA, V.P.;
GRIDINA, G.P.

Practice in dispensary treatment of bathhouse workers with
epidermophytosis. Vest. dermat. i ven. 37 no.2:70-72 P'63.

(MIRA 16:10)

1. Iz kozhno-venerologicheskikh dispanserov No.3 i 13 Leningrada.

*

ADZHIMAMUDYAN, N.I.; KEMFINSKAYA, A.V.; UZDIN, M.M.; SHILOV, M.M.;
ZAYTSEV, V.I., retsenzent; LUTOVINOV, G.V., retsenzent;
PISAREVA, Ye.I., red.

[Fundamentals of construction planning of depots and plants
for railroad transportation and of the planning of their ter-
ritories] Osnovy stroitel'nogo proektirovaniya depoi i zavodov
zheleznodorozhnogo transporta. [By] N.I.Adzhimamudian i dr.
Leningrad, Leningr. in-t inzhenerov zhel-dor. transporta in.
V.N.Obrastsova, 1963. 79 p. (MIRA 17:7)

1. Rukovoditel' gruppy Leningradskogo Gosudarstvennogo insti-
tuta proyektirovaniya na transporte (for Zaytsev). 2. Lenin-
gradskiy Gosudarstvennyy institut proyektirovaniya na transporte
(for Pisareva)

LYUBIMOV, N.N., doktor ekon. nauk, prof.; FOKIN, D.F., kand.
ekon. nauk; SHERESHEVSKIY, M.G., doktor ekon. nauk, prof.;
FISKOPPEL, F.G., doktor ekon. nauk, prof.; DYUULEN, I.I.,
kand. ekon. nauk; LOPATIN, G.S., doktor ekon. nauk, prof.;
MOGILEVCHIK, A.Ye., red.

[Foreign trade of the U.S.S.R., 1946-1963] Vneshniaia tor-
govlia SSSR (1946-1963 gg.). Pod red. I.F.Fokina. Moskva,
IMO, 1964. 189 p. (MIRA 177)

1. Moscow. Institut mezhdunarodnykh otnosheniy. 2. Kafedra
mezhdunarodnykh ekonomicheskikh otnosheniy Moskovskogo go-
sudarstvennogo instituta mezhdunarodnykh otnosheniy (for all
except Mogilevchik).

KOROBKOV, Yu.S.; PISAREVA, V.V.

Some results of investigations of E layers in the region of the Pacific Ocean. Geomag. i aer. 4 no. 5:866-872 S-O '64. (MIRA 17:11)

1. Radiofizicheskiy institut pri Gor'kovskom gosudarstvennom universitete.

L 34494-65 FBD/EWT(1)/EWG(v)/FCC/EDC-4/EEC(t)/EWA(h) Po-4/Pe-5/Pq-4/Pae-2/
Feb/PI-1 RD/GW/WS-4

ACCESSION NR: AP5005199

S/0203/65/005/001/0173/0177

AUTHOR: Korobkov, Yu. S.; Pisareva, V. V.

TITLE: Latitudinal distribution of absorption of cosmic radio emission in the ionosphere

SOURCE: Geomagnetizm i aeronomiya, v. 5, no. 1, 1965, 173-177

TOPIC TAGS: cosmic radio emission, radio wave absorption, ionospheric F2 layer, ionospheric D layer

ABSTRACT: This paper presents the results of measurements of absorption of cosmic radio emission made over the Pacific Ocean in the range of latitudes 36°N-18°S in the summer (May-August) of 1962 (between longitudes 160-240°E). Measurements were made by the two-frequency method. Fig. 1 of the Enclosure shows the mean diurnal curves of ionospheric absorption at a frequency of 22 Mc for each range of latitudes. All the diurnal curves have a characteristic maximum in the afternoon hours, and in some cases there is a small evening maximum whose nature is not clear. The latitude dependence Γ_t of total radio wave absorption in the ionosphere is shown in Fig. 2. Total absorption increases with a decrease in latitude, but in the region of the geomagnetic equator there is a noticeable maximum. The absorption

Card 1/4

L 31194-65
ACCESSION NR: AP5005199

maxima are at geographic latitudes of approximately 16°N and 11°S. The illustrated dependence of total ionospheric radio absorption on latitude indicates that there is a clearly expressed geomagnetic effect in the latitudinal distribution. For comparison, curve d in Fig. 2 shows the dependence of the critical frequency of the F2 layer for hours close to midday. Data on f_oF2 were obtained at the same time. The relationship between total absorption and f_oF2 indicates that the F2 region makes an appreciable contribution to total ionospheric absorption. On the basis of the determined latitudinal distribution of absorption of cosmic radio emission and the values f_oF2 , an attempt is made to detect the contribution Γ_F to the total absorption Γ introduced by the F2 region and the contribution Γ_D introduced by the lower regions of the ionosphere, particularly the D layer. The latitude dependence of absorption in the F2 layer Γ_F and absorption in the layer ionosphere Γ_D are shown in Fig. 2 of the Enclosure. Orig. art. has: 11 formulas, 2 figures and 1 table. [08]

ASSOCIATION: Nauchno-issledovatel'skiy radiofizicheskiy institut pri Gor'kovskom universitete (Scientific Research Institute of Radio Physics at Gorky State University)

SUBMITTED: 09Jun64

ENCL: 02

SUB CODE: ES

NO REF SOV: 005

OTHER: 011

ATD PRESS: 3213

Cord 2/4

L 34494-65

ACCESSION NR: AP5005199

ENCLOSURE: 01




Fig. 1. Mean diurnal curves of ionospheric absorption at 22 Mc for different latitude ranges

Card 3/4

L 34494-65

ACCESSION NR: AP5005199

ENCLOSURE: 02




Fig. 2. Latitude dependence of the total absorption of radio waves in the ionosphere

Card 4/4

PISAREVICH, I.N., inzhener; SLAVUTSKIY, A.K., kandidat tekhnicheskikh nauk,
nauchnyy redaktor.

[Construction of railroads and commercial highways] Stroitel'stvo
shelesnykh i avtomobil'nykh promyshlennyykh dorog. Moskva, Gos. izd-vo
lit-ry po stroitel'stvu i arkhitekture, 1954. 310 p. (MLRA 7:6)
(Railroads--Construction) (Road construction)

PISAREVICH, I. N.

Construction of railroads and commercial highways. Moskva, Gos. izd-vo lit-ry po stroitel'stvu i arkhitekture, 1954. 310 p. ma. (54-38748)


TF240.P5

1. Railroads-Construction. 2. Local construction.

Тармак, Я.І. [Tarman, Y.I.; Pysarevych, V.V.]

Effect of iprazid on the metabolism of the glutamate and protein
and protein amide groups in the adult brain. Ukr. biochem. zhurn.
35 (1987) 1-10.

1. Iprazid is a potent inhibitor of the glutamate decarboxylase
activity in the brain.



PISARNYSKAYA, Klara Isidorovna; CHUMICHEV, Aleksey Grigor'yevich;
BEREZOVSKIY, Semen Mikhailovich [deceased]; GURVITS, A.I., red.;
LANOVSKAYA, M.P., red. izd-va; BEKKER, O.G., tekhn. red.

[Operation of equipment used for the separation of scrap metal]
Eksploatatsiia oborudovaniia dlia razdelki metalicheskogo loma.
Moskva, Gos. nauchno-tekhn. izd-vo lit-ry po chernoi i tsvetnoi
metallurgii, 1958. 251 p. (MIRA 11:8)
(Scrap metal industry--Equipment and supplies)

PHASE I BOOK EXPLOITATION 955

Pisarevskaya, Klara Isidorovna; Chumichev, Aleksey Grigor'yevich; and
Berezovskiy, Semen Mikhaylovich, Deceased

Ekspluatatsiya oborudovaniya dlya razdelki metallicheskogo loma
(Operation of Equipment Used for the Preparation of Scrap Metal)
Moscow, Metallurgizdat, 1958. 251 p. 3,000 copies printed.

Ed.: Gurvits, A.I.; Ed. of Publishing House: Lanovskaya, M.R.;
Tech. Ed.: Bekker, O.G.

PURPOSE: This book is intended for skilled workers, engineers, and technicians employed at scrap-preparation depots, scrap drops, and scrap shops, as well as at plants reprocessing secondary ferrous metals. The book may also be useful to students at metallurgical tekhniums.

COVERAGE: Descriptions are given of equipment for processing iron and steel scrap, together with instructions for the operation and maintenance of the equipment, performance data, and information on

Card 1/10

Operation of Equipment (Cont.) 955

technological processes. In addition, systems of organizing scrap-preparation operations are described. The authors express their thanks to P.V.Matveyev, Engineer, for his assistance in preparing the book. There are 8 references, all Soviet.

TABLE OF CONTENTS:

Preface	7
Introduction	9
Ch. I. Breaking-up of Scrap with Skull Crackers	11
Gantry-type skull crackers	12
Tower-type skull crackers	14
Tripod skull crackers	18
Break-up devices with derrick cranes	20
Basic parts and hoisting devices	21
Foundations, anvil blocks, and pits	21
Safeguards	31
Skull-cracker balls	32
Grip tongs and hooks	33
Electric winches	37

Card 2/10

DONDE, Yu.Ya.; PISAREVSKAYA, M.A.

Using alternating current in graduating industrial optical
pyrometers. Izv.tekh. no.9:29-30 S '60. (MIRA 13:9)
(Pyrometers)

POKROVSKAYA, Taisiya Vasil'yevna; RUBINSHTEYN, Ye.S., prof., red.;
PISAREVSKAYA, V.D. red.; VLADIMIROV, O.G., tekhn.red.

[Leningrad's climate] Klimat Leningrada. Pod red. B.S. Rubinshtein.
Leningrad, Gidrometeor. izd-vo, 1957. 114 p. (MIRA 11:1)
(Leningrad--Climate)

Pisarevskaya, V. D.

PHASE I BOOK EXPLOITATION

360

Moscow. Tsentral'nyy institut prognozov

Voprosy sinopticheskoy meteorologii (Problems in Synoptic Meteorology)
Leningrad, Gidrometeoizdat, 1957. 129 p (Series: Its Trudy,
vyp. 52) 1,100 copies printed.

Sponsoring Agency: Glavnoye upravleniye gidrometeorologicheskoy
sluzhby pri Sovete Ministrov SSSR

Ed. (Title page): Tomashevich, L. V.; Ed. (inside book):
Pisarevskaya, V. D.; Tech. Ed.: Soloveychik, A. A.

PURPOSE: The collection of articles is intended for employees of
the meteorological service as well as for those interested in
the activities of the Central Institute of Forecasting.

COVERAGE: The collection of articles analyzes the causes of incorrect
short-term weather predictions and explains the nature of the
errors.

Card 1/8

Problems in Sinoptic Meteorology

360

TABLE OF
CONTENTS:

Isayeva, Ye. N. Nature of Errors in Weather Forecasting
in the Summer of 1954

3

In 1954, weather forecasting in Moskovskaya Oblast' fell short of expectations, being correct to only 73.5 percent as against a 72 percent average for the entire year. The author examines each individual cause of error and concludes that precipitation, temperature, and especially errors in forecasting the baric field of a low gradient were the deciding factors in faulty predictions. The author explains how incorrect analysis of air stratification or one of developing fronts affects the forecasting. There are 3 tables and no references.

Bachurina, A. A. Analysis of the Incorrect Weather Forecast
for May 31, 1954.

9

Card 2/8

Problems in Sinoptic Meteorology

360

The forecast for this particular date was rain at night and cool during the day. The prediction was based on the observed cyclogenesis by night (and early in the morning) on May 30. The enclosed maps show: 1) weather conditions at 3 o'clock a.m. on May 30 2) thermal and baric conditions at 6 o'clock a.m. on May 30 3) forecast for 3 o'clock a.m. for May 31 4) actual weather situation at 3 o'clock a.m. on May 31. The prediction failed: there was no rain by night and the temperature on May 31 was 22° C. The error was due to incorrect forecasting of baric pressure; this is illustrated by two additional maps. There are 5 maps and no references.

Mertsalov, A. N. Two Cases of Convective Rain

15

The article discusses two cases of erroneous weather prediction in Moskovskaya oblast' for July 29 and 30, 1954 due to convective rain. On July 28 in the evening, the prediction for the following day was no rain. This prediction was repeated the next morning. Nevertheless, it rained heavily with precipitation
Card 3/8

Problems in Synoptic Meteorology

360

mounting to 35.2 mm. The prognostics for July 30 read: scattered showers. In fact, it rained throughout the entire Moskovskaya oblast' with precipitation ranging from 8 to 18.9 mm. As a cyclone was moving westward covering the whole oblast, the rainfall was caused by convective instability. Because of an incorrect diagnosis of the baric field on the eve of the rainfall, the movement of the cyclone was not predicted in the forecast. There are 12 synoptic maps illustrating the above two cases and 3 Soviet references.

Isayeva, Ye. N. Analysis of the Erroneous Weather Forecast for July 28, 1954

31

The forecast for Moskovskaya oblast' for this date was rain. The error was caused by incorrect prediction of the movement of a cyclone approaching Moscow from the Baltic area. Two maps show the baric pressure near the surface and the thermal and baric situations on the morning of July 27. The author explains the mistake made in the analysis of this situation and shows how and why the expected cyclone by-passed Moscow. There are two synoptic maps, 1 table and no references.

Card 4/8

Problems in Synoptic Meteorology

360

Tomashevich, L. V. Analysis of the Erroneous Weather
Forecast for May 2, 1954

35

The Moscow forecast for this date, confirmed on the morning of May 2nd read: partially cloudy, no rain, with daily temperature of 20 to 22°C. The error was caused by an unexpected retardation in the movement of two warm fronts from the South, which produced rain and with it a drop in temperature to 10°C. There are 3 synoptic maps and 2 Soviet references.

Bachurina, A. A. Analysis of the Incorrect Weather Forecast
for June 26, 1954

40

The Moscow forecast for this date read: some cloudiness, no rain, daily temperature from 22 to 27°C. This was confirmed on the morning of June 26th. The error was due to incorrect evaluation of the factors causing precipitation. The capital was hit by torrential rains and the rain was persistent. Evolution of the zone of rain progressed from the direction of Card 5/8

Problems in Synoptic Meteorology

360

Smolensk but this had not been foreseen by the forecast service. There are 6 figures, 2 tables and no references.

Gorodova, M. I. Storm on July 4, 1954

47

The storm was not predicted in the morning forecast for Moscow. The synoptic map for this day was made at 3 o'clock in the morning. Although a slowly moving anticyclone was expected to reach the area of Moscow some time in the afternoon, no immediate rain was predicted. Nevertheless, the storm came at 5:30 a.m. and lasted until 11 a.m. The storm resulted from instability produced by the advection of saturated air, while the adiabatic gradient created conditions for convective rain. There are 7 drawings, 2 tables and 3 Soviet references.

Card 6/8

Problems in Sinoptic Meteorology

360

Cherkasskaya, V. M. Torrential Rains in the Ridge of High Pressure on August 12 and 13, 1954

57

For August 13th the Moscow forecast read: no precipitation. However, the whole oblast was hit in the evening by torrential rains amounting to 30 mm in the capital. The prediction was based on the position of isallohypsals lines and on the calculation of the movement of a depression, the axis of which expected to be east of Moscow towards evening. The convective instability was created by adiabatic decrease in temperature at 500 millibar level and by the advection of colder air at a 700-850 millibar level. There are 8 figures and 1 Soviet reference.

Neronova, L. M. Distribution of Summer Precipitation in Moskovskaya Oblast'

67

Since the majority of incorrect weather predictions in 1954 in Moskovskaya oblast' concerned precipitation, the author
Card 7/8

Problems in Synoptic Meteorology

360

analyzes the total distribution of rainfall throughout the entire oblast from the point of view of both intensity and occurrence. The author refers to previous attempts by I.I. Kasatkin to sum up the distribution of rainfall in the area of Moscow. The article includes a map of all meteorological stations in the oblast and draws general conclusions as to the amount of rainfall from both frontal zones and air masses. In the appendix there are tables showing maxima of precipitation under various synoptic situations (ridge, cold front, anticyclone, depression, etc.) and a listing of average monthly rainfall observed at each station. There are 9 maps, 16 tables, and 6 Soviet references in the text and 5 tables in the appendix.

AVAILABLE: Library of Congress (QC851.M64V.52)

Card 8/8

MM/vm

June 26, 1958

PHASE I BOOK EXPLOITATION

AUTHOR: See table of contents

387

TITLE: Trudy Tsentral'nogo instituta prognozov (Transactions of the Central Institute of Forecasting). Nr 51, Voprosy dolgosrochnykh prognozov (Long-term Forecast Problems)

PUB. DATA: Gidrometeorologicheskoye izdatel'stvo, Leningrad, 1957, 150 pp., 1,000 copies

ORIG. AGENCY: Glavnoye upravleniye gidrometeorologicheskoy sluzhby pri Sovete ministrov SSSR

EDITOR: Kurganskaya, V. M.; Pisarevskaya, V. D.; Tech. Ed.: Vladimirov, O. G.

PURPOSE: This collection of articles is for specialists in the field of long-term weather forecasting.

COVERAGE: The collection of articles analyzes the rhythmicity of atmospheric processes and especially those originating in polar regions, and it evaluates the possibility of using the occurrence of rhythms in weather forecasting.

Card 1/7

Transactions of the Central Institute of Forecasting (Cont.)

387

TABLE OF
CONTENTS:

- Duletova, T. A. and Komissarova, L. N. Relation Between Seasons and Rhythmicity 3
The authors refer to B. P. Mil'tanovskiy's method of long-term forecasting as standard in the USSR. They also recapitulate the basic postulate of Mil'tanovskiy's theory that natural synoptic seasons and rhythmicity in atmospheric processes are interrelated. The article expands this theory. The authors suggest making an integrated map of depressions and ridges within a certain span of time. Such a map, compiled at AT 500 (absolute topography at the 500 millibar level), would automatically reveal all deformations in atmospheric processes and their deviation from some definite synoptic patterns. There are 3 tables, 14 maps, 2 diagrams, and 3 Soviet references.

Card 2/7

Transactions of the Central Institute of Forecasting

387

Vitel's, L. A. Solar Origin of Atmospheric Rhythms

22

The author examines the relationship between solar activity and atmospheric processes and draws the following conclusions: 1. Periods of intensified solar activity can neither be ascribed to definite areas nor can they be considered constant in their degrees of intensity. 2. Although rhythmic changes in atmospheric processes are dependant on variations in solar activity, yet similar solar effects do not always produce identical responses in atmospheric rhythms. The article mentions S. T. Pagava, K. V. Brodovitskiy, P. P. Predtechenskiy, B. M. Rubashev (Pulkovo Observatory), M. N. Gnevyshev (Pulkovo Observatory), M. S. Eygenson, V. G. Shishkov, and V. V. Shuleykin as the leading scientists in the field of studies of solar impact on atmospheric processes. There are 11 diagrams, 2 maps, and 26 references, of which 20 are Soviet, 1 is French and 5 are in English.

Isayev, E. A.

Investigation of a Sharp Decline in Temperature in European USSR Caused by Certain Synoptic Processes.

The author separates the occurrence of cold waves in synoptic processes of the moderate zone of European USSR into ultra polar, meridional and normal types and remarks on the role of the advection of cold air masses from the polar region.

Card 3/7

Transactions of the Central Institute of Forecasting

387

The first chapter of the article contains general information on the nature of cold waves, and a number of anticyclonic outbreaks travelling southwards is analyzed. The existence of monthly rhythmicity in all types of processes and its application in long-term forecasts is the subject of the second part of the article. In the third part the author compiles statistical data on air temperature during the first six months of the year for Moscow, Voronezh, Penza, and Vologda and he demonstrates the probability of recurrence and rhythmicity in such repetitions. The author defines the term "sharp" decline in temperature as a decline of the average daily temperature by 5° to 10°C during cold seasons and 3° to 7°C in warm seasons provided that such temperature lapse occurs within 1-2 days. The author concludes that in addition to seasonal rhythmicity there are also monthly rhythms of synoptic processes. The statistical data are to prove that a definite successive recurrence exists among the various types of air circulation and also in the location and distribution of baric fields. Consequently, the occurrence of certain types of synoptic situations during a given period will allow the prediction of definite synoptic situations in the non-distant future. There are 11 tables, 14 maps, and 5 Soviet references.

Card 4/7

Transactions of the Central Institute of Forecasting

187

Avanesova A. G., Kaak L. I., and Yausheva G. Sh. Occurrence of Selected Ultra-polar Processes in Central Asia and Kazakhstan.

83

The authors evaluate the efficacy of long-term weather forecasts based on the periodic occurrence of ultrapolar processes. The latter are traced along their meridional extent from some definite reference points in the North, i.e., the Barents Sea, Novaya Zemlya, etc. In the appendix, 54 ultrapolar processes are analyzed and their reference localities specified. In addition, the tabular material specifies also the occurrence of respective synoptic phenomena consequent upon the appearance of polar air processes. The rhythmicity of recurrence is repeated in intervals of 3 to 5 months. There are 11 maps, 1 diagram, and 4 tables, in addition to 16 pages of tabular data in the appendix. All 7 references are Soviet.

Goncharova, Ye. F. Synoptic Conditions of the Exceptionally Cold Spring of 1952 in Northern Caucasus

117

The average daily temperature in March was 2° to 5°C below the norm and in April and May, 1° to 1.5°C. Similar conditions were observed during the springs of 1945, 1940, 1933, etc. The article analyzes these conditions. There are

Card 5/7

Transactions of the Central Institute of Forecasting

387

three main types of synoptic processes which can cause an unusually cold spring in this area: 1. The occurrence of an anticyclone in Scandinavia with a tendency to travel south. 2. The existence of a depression over the Soviet Northeast with the ensuing displacement of anticyclones towards the Caspian Sea. 3. The occurrence of a large depression over the northern half of European USSR and the constant advection of cold air into this depression. This depression has a tendency to extend as far south as the Black Sea. There are 3 maps, 1 diagram, and 1 Soviet reference.

Troshnikov M. V. Problem of Rhythmicity in Ultrapolar Synoptic Processes in Siberia and the Far East

124

The article refers to B. P. Mui'tanovskiy's contribution to the interpretation of ultrapolar processes and their behavior. Mui'tanovskiy discovered that the recurrence of conditions can be observed every 3 to 5 months. In later years S. T. Pagava proved that there are also intermediate rhythms which repeat at intervals of 45-75 days. In the present article the author not only recapitulates the work of his predecessors but also describes the nature of such polar processes. The processes are traced from some definite reference points such as the Kara Sea, Kolyma, etc. The author explains the role of these processes

Card 6/7

Transactions of the Central Institute of Forecasting

387

in synoptic forecasts and their low reliability. The appendix contains data on synoptic processes which may be similar, different, or reversed with respect to their corresponding polar processes. There are 3 maps, 5 diagrams, 8 Soviet references, 5 tables, and a 9-page appendix.

AVAILABLE: Library of Congress (QC 851.M64)

Card 7/7

GC/bmd

6 June 1958

SINEL'SHCHIKOV, V.V., otvetstvennyy redaktor; PISAREVSKAYA, V.D., redaktor;
BRAYNINA, M.I., tekhnicheskii redaktor.

[Instructions for hydrometeorological stations and posts]
Nastavlenie gidrometeorologicheskim stantsiham i postam. Lenin-
grad, Gidrometeor. izd-vo. No. 11. [Agrometeorological observations
at stations and posts] Agrometeorologicheskie nabludeniia na
stantsiakh i postakh. Pt. 1. [Principal agrometeorological
observations] Osnovnye agrometeorologicheskie nabludeniia. 1957.
203 p. (MIRA 10:11)

1. Russia (1923- U.S.S.R.) Glavnoye upravleniye gidrometeorologicheskoy
sluzhby.

(Meteorology, Agricultural)

NIKANDROV, V.Ya., kand.fiz.-mat.nauk, red.; SHISHKIN, N.S., doktor fiz.-mat.
nauk, red.; SHIPRIN, K.S., doktor fiz.-mat.nauk, red.; SOLOV'YEV,
V.A., kand.fiz.-mat.nauk, red.; PISAREVSKAYA, V.I., red.;
SOLOVYCHIK, A.A., tekhn.red.

[Investigations of clouds, precipitation, and thunderstorm
electricity] Issledovanie oblakov, osadkov i grozovogo elektri-
chestva; sbornik dokladov V Mezhdunarodnoy konferentsii po
voprosam issledovaniya oblakov, osadkov i grozovogo elektrichestva.
Leningrad, Gidrometeor. izd-vo, 1957. 214 p. (MIRA 11:6)

1. Russia (1923- U.S.S.R.) Glavnoye upravleniye gidro-
meteorologicheskoy sluzhby.

(Clouds) (Atmospheric electricity)
(Precipitation (Meteorology))

PHASE I BOOK EXPLOITATION

753

Brounov, Petr Ivanovich

Izbrannyye sochineniya. t. 1 Sinopticheskaya meteorologiya (Selected Works.
v. 1: Synoptic Meteorology) Leningrad, Gidrometeoizdat, 1957. 302 p.
1,800 copies printed.

Resp. Ed.: Vitel's, L. A., Ed.-Compiler Usmanov, R. F., Ed.: Pisarevskaya,
V.D.; Tech. Ed.: Braynina, M.I.

PURPOSE: This book is designed for the meteorologist specializing in synoptic
meteorology.

COVERAGE: In 1953 the All-Union Conference of Synoptologists resolved to publish
the selected works of P. I. Brounov (1852-1927), the founder of Russian synoptic
meteorology. Brounov, whose works had never been published in one edition
before, made important contributions in the fields of synoptic and agricultural
meteorology, atmospheric optics, and physical geography. The present edition,
Volume I, Synoptic Meteorology, contains only a part of his writings and con-
sists of 10 works published between 1879-1921. The first nine contain the
basic results of his research in synoptic meteorology; the tenth is a proposal
Card 100

Selected Works (Cont.) 753

for a reorganization of the forecasting service in Russia. The material for Volume I was selected from Brounov's earlier works. Volume I does not include material from his classic "Course in Meteorology" published in 1927, nor material from his "Temporary Barometric Maximums in Europe", which was almost wholly reproduced in a later work, "Barometric Maximums in Europe and Their Theory", and which is included in the present volume. Also included is the second part of his "Fundamentals of Weather Studies." One of Brounov's significant conclusions regarding the formation of barometric maximums was his recognition of 3 types of maximums: 1) constant maximums, 2) half yearly maximums, and 3) temporary maximums. In forming isobar maximums Brounov established that anticyclones, as well as cyclones, usually move in the direction of their greatest elongation. In examining the movement of anticyclones, he found that the center of the barometric maximum generally moves in the direction of the greatest temperature decrease. A biographical sketch and a detailed discussion of his works, theories, and experimental studies are given in the first two introductory chapters, and a complete biography of his writings at the end of the book acquaints the reader with works not included in this edition. The works are in chronological order. The text is accompanied by diagrams and tables.

Card 2/9

Selected Works (Cont.) 753

TABLE OF CONTENTS.

Foreword

Usmanov, R.F., Petr Ivanovich Brounov and His Contributions to
Synoptic Meteorology

Anomalous Distribution of Temperature in Russia in 1879

Forward Movement of Cyclones and Anticyclones in Europe and Part of
Russia

Introduction

Ch. I. Preliminary Investigation

Ch. II. Study of the Distribution of Temperature and Absolute
Humidity in Cyclones

Ch. III. A New Theory of Movement

Card 3 9

Selected Works (Cont.)

753

Ch. IV. The Relationship Between the Thermometric Gradient,
Force, and Velocity of Cyclone Movement

Ch. V. Some Outstanding Problems and Subsequent Results of Research
gations

Ch. VI. Theory of Anticyclone Movement

Ch. VII. Analysis of the Distribution of Temperature Deviation from
the Normal Around Cyclone Centers

Ch. VIII. Maps of Atmospheric Pressure Changes

Ch. IX. Relationship Between the Path of a Cyclone, the Path of
Maximum Temperature Deviation From the Normal, and the
Path of Maximum Barometric Fall

Ch. X. New Methods of Forecasting Storms

Conclusions

Card 4 9

Selected Works (Cont.) 753

Annual Rate of Temperature Deviations From the Normal in European Cyclones

European Storms and Their Prediction

Introduction

1. Synoptic maps and their compilation 104
2. Relationship between wind and the elasticity of air 105
3. Effect of the earth's rotation on moving air 106
4. Determining the earth's deflecting force 107
5. Buys Ballot's and Stephenson's laws 108
6. Barometric minimums and maximums; cyclones and anticyclones; the barometric gradient; storms 109
7. Determining the barometric gradient 110
8. Size and form of cyclones 111
9. Winds in a cyclone 112
10. The path of a particle of air in the lower parts of a cyclone; studies by Guldberg and Mohn 113
11. Surfaces of equal elasticity; ascending air currents 114
12. Phenomena accompanying cyclones 115

Card 5/9

Selected Works (Cont.)

753

13. Changes occurring in the condition of the atmosphere at a given place when a cyclone passes over. Dove's law and its explanation
14. Formation of cyclones
15. Conclusions regarding the conditions of equilibrium in air. Studies by Hann and Peslin
16. Barometer height at the center of a cyclone; forces of cyclones; their mechanical work
17. Dependence of the magnitude of the barometric gradient on the force of the wind
18. Direction and velocity of cyclone movements
19. Hypotheses of cyclone movement; Faye, Koppen, Andrews and Mohn
20. Distribution of temperature and humidity in cyclones
21. General forecasting of storms; the storm system in Russia

Barometric Maximums in Europe and Their Theory

Ch. I. Laws of Motion of Barometric Maximums

1. Three types of barometric maximums. Purpose of the investigation
2. Maps of the paths of maximums
3. Survey of maps by months
4. Direction and velocity of the forward motion of maximums

Card 6/9

Selected Works (Cont.)

753

Ch. II. Form of Barometric Maximums and Height of Barometer in the Center

5. Form of barometric maximums
6. Direction of greatest isobar elongation
7. Height of barometer in the center of maximums

Ch. III. Formation of Barometric Maximums

8. Two main systems in the formation of maximums and minimums: thermal and dynamic
9. Probable effect on the formation of maximums when cold air with a greater water vapor content is replaced by colder air with a smaller water vapor content
10. Remarks on other systems in the formation of maximums
11. Examples of thermal and dynamic formations of maximums
12. Formation of maximums in Europe, 1876-1879; maximums coming from other countries into Europe during the same period

Ch. IV. Causes of the Forward Movement of Maximums

13. Remarks on the theory of general currents
14. Temperature decrease in the area of the maximum

Card 7/9

Selected Works (Cont.)

753

15. Interrelation between the direction of greatest temperature decrease and the direction of the forward movement of the center of the maximum
16. Position of the area of greatest decrease in temperature in relation to the center of the maximum
17. Additional information on changes in temperature in the area of the maximum; summary of results
18. Effect of barometric minimums on the shift in maximums
19. Effect of exchange of lighter air by heavier air on the shift of maximums
20. Hypothesis of the forward movement of maximums
21. Examples of shift in maximums
22. Two maximums of 1885
23. Causes of different velocities in the movement of maximums
24. Cessation of maximums; their weakening and disappearance
25. Predictions of the future course of the maximum

212
212
213
222
24
27
2

Card 8'9

GAVRILOV, Valentin Aleksandrovich; PETROV, N.A., otv.red.; PISAREVSKAYA,
V.D., red.; SERGEYEV, A.N., tokhn.red.

[Transparency of the atmosphere and visibility] Prozhrachnost'
atmosfery i vidimost'. Izd.2., perer.i dop. Leningrad, Gidro-
meteor.isd-vo, 1958. 166 p. (MIRA 12:4)
(Atmospheric transparency) (Visibility)

GOL'TSBERG, I.A., doktor geogr.nauk; ARKHIPOVA, Ye.P., kand.geogr.nauk;
GLEBOVA, M.Ye.; ROMANOVA, Ye.N.; SMIRNOVA, N.V.; VORONTSOV, P.A.,
kand.fiz.-mat.nauk; BARASHKOVA, Ye.P., mladshiy nauchnyy sotrudnik;
GEDSONOV, A.D.; GOLUBOVA, T.A.; MISHCHENKO, Z.A.; FEDOSEYEV, A.P.,
kand.sel'skokhoz.nauk; BELOBORODOVA, G.G., mladshiy nauchnyy so-
trudnik; PISAREVSKAYA, V.D., red.; VOLKOV, N.V., tekhn.red.

[Microclimate of the northern part of the Kazakh hills] Mikroklimat
severnoi chasti Kazakhskogo melkosopohnika. Pod red. I.A. Gol'ts-
berg. Leningrad, Gidrometeor. izd-vo, 1958. 206 p. (MIRA 12:2)

1. Leningrad. Glavnaya geofizicheskaya observatoriya. 2. Sotrudnik
Glavnoy geofizicheskoy observatorii im. A.I. Voyeykova (for Gol'ts-
berg, Arkhipova, Glebova, Romanova, Smirnova, Vorontsov, Barashkova,
Gedsonov, Golubova, Mishchenko). 3. Sotrudnik Kazakhskogo nauchno-
issledovatel'skogo gidrometeorologicheskogo instituta (for Fedoseyev,
Beloborodova).

(Kazakhstan--Microclimatology)

PRIKHOT'KO, G.F., otv.red.; PISAREVSKAYA, V.D., red.; BRAYNINA, M.I.,
tekhn.red.

[Papers of the Conference on Agricultural Meteorology and
Climatology in the Ukrainian S.S.R.] Materialy Konferentsii po
agrometeorologii i agroklimatologii Ukrainskoi SSR. Leningrad,
Gidrometeor.izd-vo, 1958. 247 p. (MIRA 12:4)

1. Konferentsiya po agrometeorologii i agroklimatologii Ukrainskoy
SSR.
(Ukraine--Meteorology, Agricultural) (Ukraine--Crops and climate)

VENTSKEVICH, Georgiy Zenonovich; RUDENKO, A.I., otv.red.; PISAREVSKAYA, V.D.,
red.; BRAYNINA, M.I., tekhn.red.

[Agricultural meteorology] Agrometeorologiya. Leningrad, Gidro-
meteor.isd-vo, 1958. 375 p. (MIRA 12:3)
(Meteorology, Agricultural)

SOKHRINA, Reisa Fedorovna, nauchnyy sotrudnik; CHELPANOVA, Ol'ga Mikhaylovna, kand.geogr.nauk; SHAROVA, Valeriya Yakovlevna, kand.geogr.nauk. Prinimeli uchastiye: RUBINSHTAYN, Ye.S., prof.; DROZDOV, O.A., prof., doktor geograf.nauk, red.; PRIK, Z.M.; PISAREVA, G.P., nauchnyy sotrudnik; GALINA, M.B.; KOSENKOVA, Z.D.; TIKHOMIROVA, N.A.; FEDOSHEVA, G.N.. POKROVSKAYA, T.V., kand.geograf.nauk, red.; PISAREVSKAYA, V.D., red.; VOLKOV, N.V., tekhn.red.

[Air pressure, air temperature and atmospheric precipitation in the Northern Hemisphere] Davlenie vozdukha, temperatura vozdukha i atmosferye osadki severnogo polushariya. Pod red. O.A.Drozдова i T.V.Pokrovskoi. Leningrad, Gidrometeor.izd-vo, 1959. 473 p. [Atlas of charts] Atlas kart. (MIRA 13:4)
(Meteorology--Charts, diagrams, etc.)

BAGROV, N.A., red.; MORSKOY, O.I., red.; PISAREVSKAYA, V.D., red.; BRAYNINA,
M.I., tekhn.red.

[Weather forecasting; collection of translated articles] Voprosy
oredakssaniia pogody; sbornik perevodaykh statei. Leningrad,
Gidrometeor. izd-vo, 1958. 439 p. (MIRA 11:10)
(Weather forecasting)

SHAROVA, Valeriya Yakovlevna; DROZDOV, O.A., doktor geogr. nauk, red.;
PISARENKOVSKAYA, V.D., red.; SOLOVYCHIK, A.A., tekhn. red.

[Number of days with precipitation of various amounts in the
European part of the U.S.S.R. and in the Caucasus] Chislo dni
s osadkami razlichnoi velichiny na territorii Evropeiskoi chasti
SSSR i Kavkaza. Pod red. O.A. Drozdova. Leningrad, Gidrometeor.
izd-vo, 1958. 142 p. (MIRA 11:9)
(Meteorology--Charts, diagrams, etc.) (Precipitation)

GURAL'NIK, Israil' Iosifovich; MAMIKONOVA, Sof'ya Vartanovna; POLKOVNIKOV, Maksim Andreyevich; KAROL', B.P., otv.red.; PISAREVSKAYA, V.D., red.; PROTOPOPOV, V.S., red.; FLAUM, M.Ya., tekhn.red.

[Problems in meteorology] Zadachnik po meteorologii. Leningrad, Gidrometeor.izd-vo, 1959. 251 p. (MIRA 13:2)
(Meteorology--Problems, exercises, etc.)

12 11/11/11 12
BROUNOV, Petr Ivanovich; USMANOV, R.F., redaktor-sostavitel'; VITEL'S, L.A.,
otvetstvennyy red.; PISAREVSKAYA, V.D., red.; BRAYNINA, M.I., tekhn.
red.

[Selected works] Izbrannye sochineniia. Leningrad, Gidrometeor.
izd-vo. Vol.1. [Synoptic meteorology] Sinopticheskaia meteorologiya.
1957. 302 p. (MIRA 11:2)
(Meteorology)

TITOV, L.F.; SOLNTSEVA, N.O.; PISAREVSKAYA, V.D.

Calculation of fluctuations of the level of the sea during a period
of storms in the western Gulf of Finland. Trudy GOIN no.69:28-45
'62. (MIRA 15:11)

(Finland, Gulf of—Oceanography)

MIKHAYLOV, Yu.D.; PISAREVSKAYA, V.D.

Results of shipboard studies of the wind above the Gulf of Finland.
Trudy GOIN no.69:57-72 '62. (MIRA 15:11)
(Finland, Gulf of—Winds)

OPARINA, Ye.M.; SENTYURIKHINA, L.N.; PISAREVSKAYA, Ye.I.

Effect of unsaponifiables of oxidized petroleum wax on the properties of greases. Trudy VNII NP no.7:367-373 '58.

(MIRA 12:10)
(Lubrication and lubricants) (Paraffins) (Oxidation)

SEMYURIKHINA, L.N.; OPARINA, Ye.M.; PISAREVSEAYA, Ye.B.

Thickening power of calcium and lithium soaps of acids isolated
from oxidized petroleum wax. Trudy VNII NP no.7:359-366 '58.

(MIRA 12:10)

(Metallic soaps) (Paraffins)

(Lubrication and lubricants)

L 21032-66 EWP(●)/EWT(●)/EWP(J)/T WW/DJ/GS/RM/WH
ACCE3910W NR: AT5020438

UW/0000/65/000/000/0134/0136

AUTHORS: Oparina, Ye. M.; Sentyurikhina, L. N.; Dmitriyova, V. O.; Pionrovskaya, Ye. E.; Petrova, L. N.

TITLE: High temperature lubricants based on dyes

SOURCE: AN SSSR. Nauchnyy sovet po troniyu i mazkam. Teoriya mazoschnogo deystviya i novyye materialy (Theory of lubricating action and new materials). Moscow, Izd-vo Nauka, 1965, 134-138

TOPIC TAGS: lubricant, dye based lubricant, lubricant additive/ TsIATIM 221a lubricant, PFMS 4 silicone fluid, ETs 3 centrifugo, FM1322/300 silicone fluid

ABSTRACT: Lubricants based on dyes which are stable up to 350C were investigated. Polymethylphenyl-siloxane liquids with different methyl and phenyl group ratios (E.M. Oparina i dr. Khimiya i tekhnologiya topliv i masel, 1961, No. 1) were used as the dispersion media. It was found from the volumetric mechanical properties that vat dyes blue "K," indigo, dioxyviolanthrone, and dimethoxyviolanthrone have weak thickening properties while the other dyes (pigment "SA," vat dyes blue "H," "O," and isoviolanthrone) form lubricants which are similar in mechanical properties and colloidal stability to silicone lubricants (TsIATIM-221a, for example). To

L 21032-66

ACCESSION NR: AT5020430

8

determine storage stability and high temperature stability, the lubricants were tested by the KSA method (350 gm load) and on heated centrifuge ETa-3 (at 150C for 5 hours) respectively. It was found that with PFMS-4 fluid the colloidal stability of good thickening dyes was better than that of less effective thickeners and comparable to TolATIM-221a. Percent weight loss of lubricant based on different fluids (using pigment 3A) was found to be 3.0, 4.2, 6.3 and 11.0% at 250C and 0, 17.1, 18.0 and 29.1% at 300C for PFMS-4, copolymer 2/300, copolymer 3, and FM1322/300 fluids respectively. It was also found that the plastic properties, i.e., effective viscosity and strength of isoviolanthrone-based lubricants (after heat stabilization), were practically unchanged after 1000 hrs at 150C. Indanthrene and isoviolanthrone silicone lubricants were tested in ball bearings at high speeds ($D_n = 300\ 000\ \text{mm rev/min}$) at 150C and 15000 kg/cm² and were found inferior to TolATIM-221 lubricants. At lower speeds (to 10000 mm rev/min) and low loads the above lubricants operated longer than 1500 hours at 200C. Dyos can be used as thickeners in conjunction with graphite and molybdenum disulfide, giving up to 2500 hrs of service at 200C, 100 rpm, and 20000-25000 kg/cm² (lubricant NK-50 fails after 8-10 hrs under those conditions). At lesser speeds and loads service of 3000 hrs at 350C can be obtained. Orig. art. has: 4 tables.

ASSOCIATION: Nauchnyy sovet po treniya i mazaniy, AN SSSR (Scientific Committee on Friction and Lubrication, AN SSSR)

2/3

PIKAREVSKIY,
USSR/Engineering

Card 1/1 : Feb. 128—18/33

Authors : Pisarevskiy, Cand. Tech. Sci.

Title : ~~Temperature dependence of the damping of vibrations and the modulus of elasticity of some steels~~
: Temperature dependence of the damping of vibrations and the modulus of elasticity of some steels

Periodical : Vest. mash. 34/8, 61-65, Aug 1954

Abstract : An analysis is made of research conducted to ascertain the effect of heat on the damping of vibrations in machines with rapidly-moving parts. The materials involved were austenite steel, tool steel, perlite steel and high-chromium steel. Temperatures up to 1300° were used. Three Russian references: (1946-1953). Graphs; illustrations; table.

Institution :

Submitted :

PISARENKSKIY, A. A.

"Shock and the fight against it in transpleural operations on the intestine and cardia. Leningrad Pediatric Medical Inst. Hospital imeni Kuybyshev. Leningrad. 1958. (DISSERTATION For the Degree of Candidate in MEDICAL SCIENCE.)

Knizhnaya letopis'
No 33, 1958, Moscow

PISAREVSKIY, A.A.

Hemangioma of the alimentary canal. Khirurgia Supplement:18-19
'57. (MIRA 11:4)

1. Iz fekul'tetskoy khirurgicheskoy kliniki (dir. - prof. S.Z.
Gaynats) Leningradskogo Gosudarstvennogo meditsinskogo pediatriche-
skogo instituta.
(ALIMENTARY CANAL--TUMORS)

GADZHIYEV, Surkhay Aliyevich; PISAREVSKIY, A.A., red.; RULEVA, M.S.,
tekhn. red.

[Surgical treatment of mitral stenosis] Khirurgicheskoe leche-
nie mitral'nogo stenoza. Leningrad, Medgiz, 1961. 217 p.
(MITRAL VALVE--SURGERY) (MIRA 15:7)

LUKOMSKIY, G.I.; PISAREVSKIY, A.A., red.; KOKIN, N.M., tekhn. red.

[Bronchoscopy in a surgical clinic] Bronkhoskopiia v khirurgi-
cheskoi klinike. Moskva, Medgiz, 1963. 275 p.

(BRONCHOSCOPY)

(MIRA 16:3)